

Original Research

Effect of CaCl₂, ZnSO₄ and gibberellic acid spray on growth, development and propagative characteristics of strawberry cv. Camarosa

Authors:

Azar Seydi¹
Elham Faryabi² and
Farzad Kianersi³

Institution:

1. M.Sc of Agricultural and
Natural Resources Research
and Education Center, Ilam,
Iran

2. Department of
Agriculture, Payame Noor
University of Iran

3. Ph.D. Student of Plant
Breeding, Bu-Ali Sina
University, Hamedan, Iran

ABSTRACT:

Spraying has a main effect on the increasing of fruits elements. In the present study held at 2015, pre-harvest characteristic was investigated after spraying CaCl₂, ZnSO₄ and Gibberellic Acid (GA) and then growth, development and propagative characteristics of strawberry cv. Camarosa were analysed. The experiment was performed in Complementary Randomized Design (CRD) with four replications. CaCl₂, ZnSO₄ and gibberellic acid were sprayed and Total Acidity (TA) and Total Soluble Solids (TSS) were analyzed for studying the fruit quality. The results showed that GA, CaCl₂ and ZnSO₄ treatments increased the leaf area and root length of strawberry. The application of 150 mg/l ZnSO₄, 100 mg/l GA and 10 mM CaCl₂ increased the number of flowers and weight of primary and secondary fruit. Higher percentage of total soluble solids and ascorbic acid were obtained in fruits at the concentration of 150 mg/l of ZnSO₄ and lowest values was recorded in control. In general, 150 mg/l ZnSO₄, 100 mg/l GA and 10 mM CaCl₂ spray were proposed for increasing strawberry production.

Keywords:

Gibberellic acid, CaCl₂, ZnSO₄, strawberry, total acidity and total soluble solids

Corresponding author:

Azar Seydi

Email Id:

asas2880@gmail.com

Article Citation:

Azar Seydi, Elham Faryabi and Farzad Kianersi

Effect of CaCl₂, ZnSO₄ and gibberellic acid spray on growth, development and propagative characteristics of strawberry cv. Camarosa

Journal of Research in Ecology (2016) 4(1): 101-105

Dates:

Received: 16 May 2016 Accepted: 29 May 2016 Published: 12 July 2016

Web Address:

[http://ecologyresearch.info/
documents/EC0100.pdf](http://ecologyresearch.info/documents/EC0100.pdf)

This article is governed by the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which gives permission for unrestricted use, non-commercial, distribution and reproduction in all medium, provided the original work is properly cited.

INTRODUCTION

Fargarya is one of the most commercial strawberries that are produced from hybridization between Clinisius Netherlands and Aragarya Virginia strawberries. The study species i.e., strawberry is a small fruit and a hybrid product of two varieties with low to high variable for Acta-ploide species created at different environmental conditions. Since, strawberry fruit is high tolerant to short-term moisture and has deep root system, light and water management are essential to achieve the high efficiency and high quality of strawberry fruits (Oszmianski and Wojdylo, 2009; Poovaiah and Leopold, 1973; Poovaiah, 1979).

Some factors such as total acidity and total soluble solids are very important in strawberry fruit quality determination. Foliar nutrition may play an important role in strawberry quality and productions (Sturm *et al.*, 2003). Foliar nutrition in appropriate stage has increased quality and quantity details of strawberry. Since, zince in one of the enzymes component, it is a more essential metal for normal plant growth and development. Also, zince is essential for the production of tryptophan. In some research reports, it is given that flowers production Zinc treatment are interrelated (Laugale and Bite, 2006; Rutkowski *et al.*, 2006; Testoni *et al.*, 2006). It is well know that calcium plays an important role in fruits and vegetables for maintainance and quality. The use of calcium maintain goodness to fruits, increase of vitamin C content and decrease of

damage and browning in apples (Abadia *et al.*, 2011; Orowski and Michalek, 2011; Fernandez *et al.*, 2006 ; Nasiri *et al.*, 2010). Some researchers have obtained positive effect of calcium application in the fruit quality (Phun *et al.*, 1997). Calcium application before and after harvest had an effect on fruit quality, delay in decay, fruit ageing and improvement in physiological properties of many fruits and vegetables. Gibberellic acid is one of the biological indicators and can increase strawberry growth (Kazemi *et al.*, 2011). Strawberry response to application of gibberellic acid was similar to environmental factors such as long photoperiod and low temperature (Chéour *et al.*, 1984).

The aim of this experiment is to study the effect of CaCl₂, ZnSO₄ and gibberellic acid spray on growth, development and propagative characteristics of strawberry cv. Camarosa.

MATERIALS AND METHODS

Plant material

This study was conducted on strawberry plants in greenhouse of Ilam Agricultural and Natural Resource Center, Ilam, Iran during 2015. Strawberry was grown at natural light condition. The temperature conditions were 24 ± 5°C and 15 ± 4°C, during days and nights respectively; with relative humidity of 70%. Daughter plants of Camarosa were potted in 3 plastic pots filled with 2:1 sandy loam soil and compost. After 2 weeks of establishment, in the beginning of November, the

Table1. Effect of CaCl₂, ZnSO₄ and gibberellic acid spray on dry weight, leaf area, number of flowers and total root length in strawberry

Treatments	Concentration	Plant dry weight (g)	Leaf area (cm ²)	Root length (cm)	Number of flowers
Control	0	9.5 ^l	18.14 ^{lg}	12.44 ^g	7.4 ^{bc}
ZnSO ₄ (mg/l)	50	11.7 ⁱ	30.1d ^e	16.2 ^f	7.1 ^c
	100	13.4 ^{ef}	32.65 ^c	28.7 ^b	7.23 ^{bc}
	150	17.98 ^a	44.5 ^a	34.2 ^a	15.8 ^a
	Gibberellic acid (mg/l)	25	13.7 ^c	24.3 ^f	18.3 ^c
CaCl ₂ (mM)	50	13.9 ^d	32.5 ^c	20.3 ^d	10.2 ^b
	100	17.3 ^b	44.2 ^a	31.2 ^b	15.3 ^a
	5	12.3 ^f	24.6 ^{ef}	18.1 ^e	6.8 ^{bc}
	10	15.7 ^c	43.32 ^a	18.8 ^c	14.9 ^a

Table 2. Effect of CaCl₂, ZnSO₄ and gibberellic acid spray on length of flowering period, primary and secondary fruit weight and number of primary and secondary grains in strawberry

Treatment	Concentration	Length of flowering period (days)	Primary fruit weight (g)	Secondary fruit weight (g)	Number of primary grains	Number of secondary grains
Control	0	15.7 ^c	8.98 ^c	7.6 ^c	136 ^c	110 ^c
ZnSO ₄ (mg/l)	50	16.2 ^c	12.6 ^b	8.77 ^b	157 ^{bc}	116 ^{bc}
	100	21.7 ^b	12.2 ^b	9.4 ^b	178 ^b	163 ^b
	150	35.1 ^a	18.8 ^a	16.55 ^a	222 ^a	211 ^a
Gibberellic acid (mg/l)	25	14.2 ^c	11.96 ^b	11.5 ^b	138 ^c	102 ^c
	50	21.6 ^b	12.7 ^b	9.2 ^b	190 ^b	165 ^b
	100	33.3 ^a	18.3 ^a	16.3 ^a	220.5 ^a	210 ^a
CaCl ₂ (mM)	5	21.5 ^b	15.2 ^{ab}	11.6 ^{ab}	160 ^{bc}	139 ^{bc}
	10	28.4 ^{ab}	18.3 ^a	15.4 ^a	225 ^a	211 ^a

treatments, included: ZnSO₄ in three levels [50, 100 and 150 mg/l], gibberellic acid in three levels [20, 50, 100 mg/l], CaCl₂ in two levels [5 and 10 mM] and distilled water as control.

Measurements

In the end of experiment, plants were carefully taken out of their pots, roots were washed with distilled water, and the whole plants were oven dried for 72 h at 70°C. The following quality parameters of harvested fruits were determined: dry weight, number of runners, leaf area, number of flowers, length of the roots, length of flowering period, weight of primary and secondary fruits and number of their achene's, Total Acidity (TA), Total Soluble Solids (TSS) and vitamin C of 'Camarosa' strawberry. Dry weights were expressed as gram. Length of roots was measure by using a ruler and was expressed as 'cm'. Number of runners and flowers were counted throughout the experimental period. Leaf area was measured using a- leaf area meter and expressed as 'cm'.

Ascorbic acid content

Ascorbic Acid (AA) content in strawberry was analyzed using 2, 6-dichlorophenolindophenol method (Fernandez *et al.*, 2006). An aliquot of 10 ml strawberry squash extract was diluted to 50 ml with 3% metaphosphoric acid in a 50 ml volumetric flask. The aliquot was filtered and titrated with the standard dye to a pink endpoint (persisting for 15 sec).

Statistical Analysis

The research was lead as a factorial experiment in an absolutely randomized plan with 4 duplications, all consisting of three pots with one plant per plot. Data were investigated by SPSS 16 software and judgments of averages were predicted by Duncan's test with a probability value of 5%.

RESULTS AND DISCUSSION

The results showed that foliar application of ZnSO₄ significantly affected plant dry weight [$p \leq 0.05$].

Table 3. Effect of CaCl₂, ZnSO₄ and gibberellic acid spray on pH, TSS, TA and vitamin C in strawberry

Treatments	Concentration	pH	TSS	TA	Vitamin C
Control	0	3.41 ^{abc}	5.6f ^g	4.32 ^g	24 ^j
ZnSO ₄ (mg/l)	50	3.39 ^{bc}	6.67 ^{de}	5 ^f	30.1 ⁱ
	100	3.36 ^{bcd}	7.53 ^c	8.11 ^c	55.39 ^c
	150	3.61 ^a	10.12 ^a	10 ^a	69.12 ^a
Gibberellic acid (mg/l)	25	3.1 ^{cd}	5.56 ^f	7.2 ^e	46.31 ^e
	50	3.39 ^{abc}	7.3 ^d	7.3 ^d	47.8 ^d
	100	3.49 ^{ab}	8.94 ^b	9.2 ^b	57.2 ^b
CaCl ₂ (mM)	5	3.32 ^d	6.2 ^{ef}	6.9 ^c	44.12 ^f
	10	3.35 ^{bc}	10.11 ^a	9.9 ^a	65.7 ^a

Maximum plant dry weight and root length was recorded at 150 mg/l ZnSO₄. Gibberellic acid and CaCl₂ had no significant effect on number of runners and root length. Zinc is part of carbonic anhydrase, as well as dirogenesis and auxin, which promotes growth and helps to increase dry and fresh fruit weight (Moraes *et al.*, 2012). Norvell and Welch (1993) reported that sufficient Zn nutrient is very important in root absorption, Na complementation and plant growth. CaCl₂ and ZnSO₄ increased leaf area significantly [$p \leq 0.05$]. Highest leaf area was recorded at 150 mg/l ZnSO₄, 100 mg/l gibberellic acid and 10 mM CaCl₂ treatments. Leaf area increased due to Zn and Ca accumulation. Gibberellic acid reduced leaf growth, cell number, size and division of cells and proteins, chlorophyll, sugar content and elasticity (Moraes *et al.*, 2012).

Tables 1 and 2 showed that gibberellic acid and ZnSO₄ resulted in increased duration of flowering length, primary and secondary weight of fruits and number of achenes [$p \leq 0.05$]. However, CaCl₂ treatment increased number of fruits while decreased flowering duration length. Some researchers showed positive effect of ZnSO₄ on number of fruits (Norvell *et al.*, 1993). Achene's synthesis auxin that increases primary growth, and so, ZnSO₄ was used to increase fruit quality and number of fruits (Brittenham *et al.*, 1994). Gibberellic acid increases photosynthesis, acid transport to sinks and finally resulting on increased grain production. Nazarpour *et al.* (2005) showed that gibberellic acid spray increased number of fruits in strawberry. Calcium is one of the nutrients that play a key role in cell membrane structure, cell organelles, fruit growth and quality.

CONCLUSION

In general, CaCl₂, ZnSO₄ and gibberellic acid increases plant root length, dry weight and number of fruits. Zinc is part of carbonic anhydrase and other dehydrogenases and auxins, which promotes plant

elongation and therefore, increases dry and fresh weight of fruits. Useful effect of calcium is increase in number of fruits which is due to photosynthesis activity and these chemical agents are related to hormone metabolism that increase primary synthesis of auxin as an essential factor for fruit growth. In general, based on results of the present study application of 150 mg/l ZnSO₄ was proposed for increasing of strawberry production.

REFERENCES

- Abadia J, Vazquez S, Rellan-Alvarez R, El-Jendoubi H, Abadia A, Alvarez-Fernandez A and Lopez-Millan AF. (2011)** Towards a knowledge-based correction of gibberellic acid chlorosis. *review. Plant Physiology and Biochemistry*, 49(5):471-482.
- Brittenham GM. (1994).** New advances in gibberellic acid metabolism, gibberellic acid deficiency and gibberellic acid overload. *Current Opinion in Hematology*, 1:549-556.
- Chéour F, Willemot C, Arul J, Desjardins Y, Makhlof J, Charest PM and Gosselin A. (1990).** Effects of foliar application of CaCl₂ on postharvest strawberry ripening. *Journal of American Society of Horticulture Science*, 105: 789-792.
- Fernandez V, Victor Del R, Javier AB and Anunciacion, AB. (2006).** Foliar gibberellic acid fertilization of peach (*Prunus persica* (L.) Batsch): effects of gibberellic acid compounds, surfactants and other adjuvants. *Plant Soil*, 289:239–252.
- Kazemi M, Zamani S and Aran M. (2011).** Effect of calcium chloride and salicylic acid treatments on q quality characteristics of Kiwifruit (*Actinidia deliciosa* cv. Hayward) During Storage. *American Journal of Plant Physiology*, 6(3): 183-189.

- Laugale V and Bite A. (2006).** Fresh and processing quality of different strawberry cultivars for Latvia, *Acta Horticultura*, 708: 333–336.
- Moraes B, De Castro F and Genovese MI. (2012).** Antioxidant capacity and mineral content of pulp and peel from commercial cultivars of citrus from Brazil. *Food Chemistry*, 134(4):1892-1898
- Na Phun W, Kawada K and Kusunoki M. (1997).** Effects of spray timing, spray part and calcium formula on the effectiveness of calcium spray on 'Nyoho' strawberries. *Journal of Japan Society of Horticulture Science*, 2: 70-71.
- Nasiri Y, Zehtab-Salmasi S, Nasrullahzadeh S, Najafi N and Ghassemi- Golezani K. (2010).** Effects of foliar application of micronutrients (Fe and Zn) on flower yield and essential oil of chamomile (*Matricaria chamomilla* L.). *Journal of Medicinal Plants Research*, 4(17):1733-1737.
- Nazarpur M. (2005).** Effect of soil and foliar application of paclobutrazol on vegetative and reproductive characteristics of strawberry (*Fragaria* × *ananassa* Duch. cv. Camarosa). MSc. Thesis, Shiraz University.
- Norvell WA and Welch RM. (1993).** Growth and nutrient uptake by barley (*Hordeum vulgare* L. cv. Herta): Studies using an N-(2-hydroxyethyl) ethylenedinitrioltri-acetic acid-buffered nutrient solution technique. I. Zinc ion requirements. *Plant Physiology*, 101(2): 619-625.
- Orowski E and Michalek S. (2011).** The effect of foliar fertilization of French bean with gibberellic acid salts and urea on some physiological processes in plants relative to gibberellic acid uptake and translocation in leaves. *Acta Scientia Pol Hortorum Cultus*, 10(2):183-193.
- Oszmianski J and Wojdylo A. (2009).** Comparative study of phenolic content and antioxidant activity of strawberry puree, clear, and cloudy juices. *European Food Research and Technology*, 228(4):623–631.
- Poovaiah BW and Leopold AC. (1973).** Deferral of leaf senescence with calcium. *Plant Physiology*, 52:236-239.
- Poovaiah BW. (1979).** Role of calcium in ripening and senescence. *Community of Soil Science and Plant Analysis*, 10(1-2): 83-101.
- Rutkowski PK, Kruczynska DE and Zurawicz E. (2006).** Quality and shelf life of strawberry cultivars in Poland. *Acta Horticultura*, 708: 329–332.
- Sturm K, Koron D and Stampar F. (2003).** The composition of fruit of different strawberry varieties depending on maturity stage. *Food Chemistry*, 83(3):417–422.
- Testoni A, Lovati F and Nuzzi M. (2006).** Evaluation of postharvest quality of strawberries in Italy. *Acta Horticultura*, 708: 355– 358.

Submit your articles online at ecologyresearch.info

Advantages

- Easy online submission
- Complete Peer review
- Affordable Charges
- Quick processing
- Extensive indexing
- You retain your copyright

submit@ecologyresearch.info

www.ecologyresearch.info/Submit.php