

Original Research

Modeling of leaf area of three Afromontane forest tree species through linear measurements

Authors:

Titus Fondo Ambebe¹,
Forba Gillian Zee² and
Martin Ambe Shu²

Institution:

1. Department of Crop
Production Technology,
College of Technology, The
University of Bamenda, P.O.
Box 39 Bambili, Cameroon.

2. Department of
Environment and Natural
Resource Management, Pan
African Institute for
Development – West Africa,
P.O. Box 133 Buea,
Cameroon.

Corresponding author:

Titus Fondo Ambebe

ABSTRACT:

Leaf area is an important parameter for evaluating growth, development and competitive ability of forest trees. Destructive methods of leaf area determination often require substantial financial investment and technical knowledge, both of which are unavailable in some developing countries and most parts of others. The aim of this study was to develop linear regression models for the non-destructive determination of leaf area of three Afromontane forest tree species (*Cordia millenii* Bak., *Gmelina arborea* Roxb. and *Entandrophragma angolense* (Welw.) C.DC.). Leaf samples were collected from trees in the Tubah Upland Forest in the Bamenda Highlands of Cameroon. Leaf length (L), maximum leaf width (W), and leaf area (LA) were determined. Linear regression analysis was conducted between LA as dependent variable and L, W, LW, L², W², L²W, LW² as independent variable and F-test was used to test for significance of the model. Goodness-of-fit was evaluated from the coefficient of determination (r^2) and Root Mean Square Error (RMSE). Values of predicted leaf area were then plotted against those of the observed leaf area. The findings showed that the leaf area of *C. millenii*, *G. arborea* and *E. angolense* can be reliably determined from the models $LA = 471.59 + 0.10LW^2$ ($r^2 = 0.85$, RMSE = 170.91, $p = 0.00$), $LA = -972.63 + 110.90W$ ($r^2 = 0.92$, RMSE = 102.72, $p = 0.00$), and $LA = -31.81 + 2.88LW$ ($r^2 = 0.97$, RMSE = 66.00, $p = 0.00$), respectively. Further studies involving additional formulae derived from L and W are, however, encouraged.

Keywords:

Leaf size, Leaf dimensions, Montane forest, Linear regression.

Article Citation:

Titus Fondo Ambebe, Forba Gillian Zee and Martin Ambe Shu

Modeling of leaf area of three Afromontane forest tree species through linear measurements

Journal of Research in Ecology (2018) 6(2): 2434-2441

Dates:

Received: 05 Sep 2018

Accepted: 21 Sep 2018

Published: 08 Oct 2018

Web Address:

[http://ecologyresearch.info/
documents/EC0629.pdf](http://ecologyresearch.info/documents/EC0629.pdf)

This article is governed by the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which gives permission for unrestricted use, non-commercial, distribution and reproduction in all medium, provided the original work is properly cited.

INTRODUCTION

The Cameroon highlands ecoregion comprises of highland forests and grassland patches mainly above 900 m elevation scattered along the border area between Cameroon and Nigeria (WWF, 2017). Along an altitudinal gradient, the vegetation changes from submontane to montane forests and ultimately subalpine grassland. Apart from having one of the highest levels of endemism among vascular plants in Africa (Ndenecho, 2011), the forests provide a wide range of ecosystem services. For instance, they constitute a vital source of medicine, food, wildlife forage, building material, and raw materials for rural artisan industries. In addition, the highlands form essential water catchments. However, most of the forest that once totally covered the Bamenda Highlands component of the ecoregion has been cleared. In fact, more than 50% of the forest cover has been lost since the 1960s (Ernest Vunan, pers. comm.). The forests are found on fertile volcanic soils, which in combination with adequate rainfall, have attracted a lot of farmers. The high human population density in the region has subjected the remaining forest to enormous pressure of cultivation and grazing. The degradation of the forests by humans is likely to be accentuated by changing global climatic conditions. The protection of the forest remnants, development of adaptive strategies to climate change, and restoration of degraded portions of the ecosystem are imperative for a sustainable supply of the benefits that are contained in the Bamenda Highlands biome.

Leaf area is an important determinant of plant performance in open and closed forests. The parameter is indicative of light interception, growth (Ghoreishi *et al.*, 2012) and plant competitive ability in the field (Zadeh *et al.*, 2011). It is an essential component of plant growth analysis and often used in plant ecophysiology research (Lambers *et al.*, 2008). Leaf area is routinely measured in experiments where some physiological phenomena like photosynthesis, respiration, water

and nutrient uptake, and transpiration are of interest (Calvo-Alvardo *et al.*, 2008). Several techniques have been presented for leaf area determination which can, however, be classified as either destructive or non-destructive (Marshall, 1968). Destructive methods are those that require the harvesting of leaves at the time of evaluation. They are performed with the use of automated leaf area meters, planimeters, scanners, and image processing software (Easlon and Bloom, 2014) among others. However, some of the equipment require substantial financial investment and a high level of technical competence for operation and maintenance (Ugese *et al.*, 2008), both of which may not be readily available in some developing countries and most parts of others. Non-destructive techniques do not require defoliation and are based on regression models that estimate leaf area from linear measurements of leaf length (L), maximum leaf width (W), L^2 , W^2 or various combinations of these variables (Chanda and Singh, 2002; Çirak *et al.*, 2005; Ilkae *et al.*, 2011; Ogoke *et al.*, 2015). The leaf area estimation models can provide researchers with many advantages. First, they constitute an alternative to the high cost and complexity associated with destructive techniques while also maintaining a high level of accuracy (Ghoreishi *et al.*, 2012). Second, leaf area measurements can be carried out on the same plants for the entire duration of a study (José *et al.*, 2014), minimizing experimental variability (De Swart *et al.*, 2004). Though fast and reliable, dealing with a large number of leaves can result in destructive sampling techniques being more time-consuming and laborious than using mathematical models for leaf area determination (Mokhtarpour *et al.*, 2010; Ilkae *et al.*, 2011).

Cordia millenii Bak. (Boraginaceae), *Gmelina arborea* Roxb. (Verbanaceae), and *Entandrophragma angolense* (Welw.) C.DC. (Meliaceae) are tree species of great ecological and economic importance (Adam and Krampah, 2005; Tchinda, 2008; Jiofack, 2010) in the Bamenda highlands that have experienced notable

declines in population. For instance, only *ca* 200 acres are left of the hitherto *ca* 400 acres of Tubah Upland forest (Ndimuh, 2017) that constituted a major habitat for the trees within the Bamenda highlands. There is a general lack of automated and digital equipment for measuring leaf area in the region. Given that the trait is quite valuable in evaluating plant acclimation to field conditions (Lambers *et al.*, 2008), its determination from regression equations is crucial for the putting in place of suitable management practices that ensure competitive advantage on regeneration sites and optimize growth of trees in plantations and natural forests. To the best of our knowledge, there are no published works on leaf area prediction models for the species. Consequently, this study was aimed at developing regression equations for the non-destructive estimation of leaf area of *C. millenii*, *G. arborea*, and *E. angolense* from leaf linear measurements.

MATERIALS AND METHODS

Study setting

The site for sample collection was the Tubah Upland Forest of the Bamenda highlands. Tubah Sub-division (4°50' – 5°20'N, 10°35' – 11°59'E; 950 – 1500 m a.s.l.) is located some 15 km from Bamenda, the capital city of the North West Region of Cameroon. The Sub-division covers four main villages; Bambili, Bambui, Kedjom-Keku, and Kedjom-Ketinguh (Nguh and

Maluh, 2017). The vegetation is mainly Savannah grassland. Its forested area is found at the northern part, mainly Kedjom-Keku where the samples in this study were obtained. A high population density and unsustainable human behavior related to farming and grazing have largely destroyed the forest and grassland vegetation (Melle *et al.*, 2016). Tubah is characterized by a dry (November to April) and a rainy season (May to October). Average annual rainfall varies from 1780 to 2290 mm and most of the precipitation occurs between July and September (Ndenecho, 2010). Average annual temperature is 20.67°C (Yuninui, 1990), with a 20-22°C / 13-14°C maximum / minimum (Kiteh, 2011). Relative humidity is highest during the months of July and August (> 80%) and lowest in January and February (45 to 52%) (Ndenecho, 2010; Ndenecho, 2011).

Study design and data collection

The donors consisted of five trees each of *C. millenii*, *G. arborea*, and *E. angolense*. To capture as much morphological variability as possible for sampling, the vertical axis of the canopy was partitioned into four fractions of approximately similar height. A fully expanded leaf was collected from each of the fractions, resulting in twenty leaves per tree species. Only leaves that had their edges preserved and had not suffered herbivore or insect damage were harvested. After the collection, the leaf was immediately sealed in an airtight polythene bag. All the samples from each tree spe-

Table 1. ANOVA table for regression model

S. No	Source	df	SS	MS	F	p-value
1	Regression	1	SSR	$MSR = \frac{SSR}{1}$	$\frac{MSR}{MSE}$	$\Pr\left(F > \frac{MSR}{MSE}\right); F \sim F_{1,n-2}$
2	Error	n-2	SSE	$MSE = \frac{SSE}{n-2}$		
3	Total	n-1	SST			

Table 2. Results of Pearson correlation between observed leaf area and variables from leaf linear measurements

S. No	Species	Variable	L	W	LW	L ²	W ²	L ² W	LW ²
1	<i>C. millenii</i>	LA	0.79	0.83	0.92	0.77	0.81	0.86	0.92
2	<i>G. arborea</i>	LA	0.94	0.96	0.95	0.93	0.96	0.93	0.95
3	<i>E. angolense</i>	LA	0.96	0.95	0.99	0.96	0.96	0.97	0.98

cies were subsequently bulked in a large leak-proof polythene bag and transported to the Pan African Institute for Development – West Africa Support Centre in Bamenda for measurement of leaf area and linear dimensions.

Observed leaf area was obtained with the use of a graph paper. The graph paper consisted of 1 cm² cells each subdivided into twenty-five 0.04 cm² cells. The leaf was fully spread out on the graph sheet and its margin traced. The observed leaf area was determined from the number of 1 cm² and 0.04 cm² cells that fell within the boundary of traced region. Leaf length (L: distance from lamina tip to lamina-petiole junction along the midrib) and maximum leaf width (W: widest distance across lamina perpendicular to length) were measured.

Data analysis

The data for each tree species were subjected to Pearson-Moment correlation for determination of the

relationship between observed leaf area and leaf linear dimensions (L, W), their squares and products (LW, L², W², L²W, LW²). The data were then subjected to linear regression analysis using the observed leaf area as dependent variable and L, W, LW, L², W², L²W, and LW² as independent variables. Statistical significance of the regression models was determined from *F*-test (see Table 1). Goodness-of-fit was evaluated from the coefficient of determination (*r*²) and Root Mean Square Error (RMSE). Selection of the suitable model for leaf area estimation was based on highest *r*² and least RMSE.

$$\text{SSR} = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$$

$$\text{SSE} = \sum_{i=1}^n (y_i - \hat{y}_i)^2$$

$$\text{SST} = \sum_{i=1}^n (y_i - \bar{y})^2$$

Table 3. Regression models of relationships between observed leaf area and variables from leaf linear measurements

S. No	Species	Variable	Model	<i>r</i> ²	<i>p</i>	RMSE
1	<i>C. millenii</i>	L	LA = -255.54 + 67.01L	0.62	0.00	268.82
		W	LA = -1059.31 + 128.78W	0.67	0.00	243.76
		LW	LA = 98.83 + 2.72LW	0.84	0.00	176.94
		L ²	LA = 594.37 + 1.27L ²	0.60	0.00	277.79
		W ²	LA = 231.63 + 3.14W ²	0.66	0.00	254.38
		L ² W	LA = 650.69 + 0.06L ² W	0.74	0.00	221.41
		LW ²	LA = 471.59 + 0.10LW ²	0.85	0.00	170.91
2	<i>G. arborea</i>	L	LA = -401.09 + 66.82L	0.89	0.00	119.54
		W	LA = -972.63 + 110.90W	0.92	0.00	102.72
		LW	LA = 162.52 + 2.17LW	0.91	0.00	109.14
		L ²	LA = 304.65 + 1.50L ²	0.87	0.00	130.21
		W ²	LA = -3.94 + 3.09W ²	0.92	0.00	102.79
		L ² W	LA = 480.96 + 0.06L ² W	0.87	0.00	131.95
		LW ²	LA = 480.42 + 0.08LW ²	0.90	0.00	118.34
3	<i>E. angolense</i>	L	LA = -610.43 + 59.29L	0.93	0.00	111.68
		W	LA = -1230.47 + 175.18W	0.90	0.00	127.42
		LW	LA = -31.81 + 2.88LW	0.97	0.00	66.00
		L ²	LA = 168.84 + 1.07L ²	0.91	0.00	121.44
		W ²	LA = -183.15 + 7.10W ²	0.92	0.00	116.46
		L ² W	LA = 341.54 + (-0.06)L ² W	0.94	0.00	102.99
		LW ²	LA = 242.83 + 0.16LW ²	0.97	0.00	75.36

where, \hat{y} is predicted leaf area; y and \bar{y} are observed leaf area and the mean, respectively.

A regression plot was established between values of predicted leaf area and those of observed leaf area for the appropriate model for each of the tree species.

RESULTS AND DISCUSSION

The value of Pearson-Moment correlation coefficient (r) was positive for the relationship between observed leaf area and each leaf linear trait (L, W, their squares and products) for the three tree species (Table 2). Relationships of the sort have been established pre-

viously for neotropical rainforest trees (Brito-Rocha *et al.*, 2016), vegetable (Schwarz and Karling, 2001) and fruit (Ramkhelawan and Brathwaite, 1990) crops. However, the leaf trait(s) that showed the strongest correlation to the observed leaf area was inconsistent across species, suggesting that the leaf area of the tree species cannot be estimated with the use of a common parameter or regression model. There was, however, a general variability in r among the tree species (Table 2), reflecting phylogenetic differences in leaf shapes.

Non-destructive estimations of leaf area from leaf dimensions are well documented. The literature presents different types of regression models that are




Figure 1. Relationship between predicted and observed leaf areas of *Cordia millenii* Bak. (a), *Gmelina arborea* Roxb. (b), and *Entandrophragma angolense* C.DC. (c).

used in the process, including power (e.g. Misle *et al.*, 2013), second-order polynomial (e.g. Mazzini *et al.*, 2010) and linear (e.g. Ogoke *et al.*, 2015) models. In this study, linear regression models were adopted because they are simpler than non-linear regression models and have yielded reliable results for a great number of plant species. The linear relationships between observed leaf area and linear functions of L or W, their squares and products were highly significant for each species (Table 3). Overall, 60 - 97% of variations in leaf area were explained by the leaf linear measurements and their derivatives. The regression with LW^2 had the highest r^2 for *C. millenii*, suggesting that the leaf area of this tree species can be estimated with the use of the model $LA = 471.59 + 0.10LW^2$ (Table 3). The selection of the appropriate model for *G. arborea* and *E. angolense* was not as direct because two equations coincided in yielding the highest r^2 (Table 3). The independent variables in the latter equations were W and W^2 for *G. arborea* and LW and LW^2 for *E. angolense* (Table 3). However, the models involving W and LW had lower RMSE, highlighting $LA = -972.63 + 110.90W$ and $LA = -31.81 + 2.88LW$ as the models for a reliable and non-destructive estimation of leaf area of the species (Table 3). The three linear regression models explained 85%, 92%, and 97% of variations in leaf area in *C. millenii*, *G. arborea*, and *E. angolense*, respectively. The finding that a single linear measurement can be used for predicting *G. arborea* leaf area is in agreement with the works of Flavio and Marcos (2003) and Ogoke *et al.* (2015). According to Brito-Rocha *et al.* (2016), the use of a single measurement (L or W) could save up to half of the time that is required for the measurement of both L and W.

Regression analysis and F-test showed a highly significant ($P = 0.00$) relationship between observed and predicted leaf areas for all three tree species. The high positive values of r^2 associated with the latter models (Figure 1) lend credence to the conclusion that the linear

models developed in this study can provide reliable estimates of leaf area of *C. millenii*, *G. arborea*, and *E. angolense*.

CONCLUSION

Linear regression models have been developed for predicting leaf area of *Cordia millenii* Bak., *Gmelina arborea* Roxb., and *Entandrophragma angolense* (Welw.) C.DC. The functions to be put into the equations can be obtained by simple measurement with a ruler. It is, hence, possible to determine the leaf area of these important tree species in a cheap and practical way. The findings of this research can have important implications for forest management given the salient role that leaf area measurement plays in assessing physiological fitness and competitive ability of plants in plantations and natural forests. However, only seven independent variables were tested in this study. Since several other combinations of L and W are mathematically possible, further studies with additional formulae are encouraged.

ACKNOWLEDGEMENT

We thank Mr. Ernest Vunan for helping us access the Tubah Upland Forest in Kedjom-Keku and assisting in the collection of samples.

REFERENCES

- Adam KA and Krampah E. 2005. *Gmelina arborea* Roxb. Ex Sm. In: In: Louppe, D., Oteng-Amoako AA and Brink M (Editors). PROTA (Plant Resources of Tropical Africa/Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands; [cited 2018 September 1]. Available from: <http://prota4u.org/search.asp>.
- Brito-Rocha E, Schilling AC, Dos Anjos L, Piotto D, Dalmolin AC and Mielke MS. 2016. Regression models for estimating leaf area of seedlings and adult individuals of Neotropical rainforest tree species. *Brazilian*

Journal Biology, 76(4): 983-989.

Calvo-Alvarado JC, McDowell NG and Waring RH. 2008. Allometric relationships predicting foliar biomass and leaf area:sapwood area ratio from tree height in five Costa Rican rain forest species. *Tree Physiology*, 28 (11): 1601-1608.

Chanda SV and Singh YD. 2002. Estimation of leaf area in wheat using linear measurements. *Plant Breeding and Seed Science*, 46(2): 75-79.

Çirak C, Odabas MS, Sağlam B and Ayan AK. 2005. Relationship between leaf area and dimensions of selected medicinal plants. *Research in Agricultural Engineering*, 51(1): 13-19.

De Swart EAM, Groenwold R, Kanne HJ, Stam P, Marcelis LFM and Voorrips RE. 2004. Non-destructive estimation of leaf area for different plant ages and accessions of *Capsicum annum* L. *The Journal of Horticultural Science and Biotechnology*, 79(5): 764-770.

Easlon MH and Bloom AJ. 2014. Easy Leaf Area: Automated digital image analysis for rapid and accurate measurement of leaf area. *Application in Plant Sciences*, 2(7): 1400033.

Flavio FB and Marcos BF. 2003. A new method for estimating the leaf area index of cucumber and tomato plants. *Horticultura Brasileira*, 21(4): 666-669.

Ghoreishi M, Hossini Y and Maftoon M. 2012. Simple models for predicting leaf area of mango (*Mangifera indica* L.). *Journal of Biology and Earth Science*, 2(2): B45-B53.

Ilkaee MN, Paknejad F and Zavareh M. 2011. Prediction model of leaf area in soybean (*Glycine max* L.). *American Journal of Agricultural and Biological Sciences*, 6(1): 110-113.

Jiofack TRB. 2010. *Cordia millenii* Baker. [Internet] Record from PROTA4U. Lemmens RHMJ, Louppe D and Oteng-Amoako AA (Editors). PROTA (Plant Resources of Topical Africa/Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands; [cited 2018 September 1]. Available from: <http://prota4u.org/search.asp>.

José JV, Fernandes RDM, Marques PAA, Ferreira ALL, Francisco JP and Duarte SN. 2014. Basil leaf area by allometric relations. *Journal of Medicinal Plant Research*, 8(43): 1275-1283.

Kiteh CC. 2011. Stakeholders participation: Myth or reality? A case study of community water supply management in Bambui-Tubah village in Cameroon. Master Thesis. Institutionen för geovetenskaper, Uppsala Universitet, Sweden. 79 p.

Lambers H, Chapin FS III and Pons TL. 2008. Plant physiological ecology. Springer, New York. 604 p.

Marshall JK. 1968. Methods of leaf measurement of large and small leaf samples. *Photosynthetica*, 2: 41-47.

Mazzini RB, Ribeiro RV and Pio RM. 2010. A simple and non-destructive model for individual leaf area estimation in citrus. *Fruits*, 65(5): 269-275.

Melle EM, Nkwatoh AF, Achiri AB and Hele TN. 2016. The unsustainable exploitation of wildlife-habitat resources in Tubah Sub-division, North West Region, Cameroon. *International Journal of Current Research in Biosciences and Plant Biology*, 3(8): 32-38.

Misle E, Kahlaoui B, Hachicha M and Alvarado P. 2013. Leaf area estimation in muskmelon by allometry. *Photosynthetica*, 51(4): 613-620.

Mokhtarpour H, Teh CBS, Saleh G, Selamat AB, Asadi ME and Kamkar B. 2010. Non-destructive estimation of maize leaf area, fresh weight, and dry weight using leaf length and leaf width. *Communications in*

Biometry and Crop Science, 5(1): 19-26.

Ndenecho EN. 2010. Implications of fuel wood yield, availability and harvest in Tubah Mountain Forest, Cameroon. *Global Journal of Human Social Science*, 10 (3): 49-55.

Ndenecho EN. 2011. Local livelihoods and protected area management: Biodiversity conservation problems in Cameroon. Langaa RPCIG, Bamenda, Cameroon. 218 p.

Ndimuh BS. 2017. Abongfen forest in real big trouble: May disappear in five years! Shanchi Ndimuh's Blog [cited 2018 August 20] Available from: Bshanchi.wordpress.com.

Nguh BS and Maluh NB. 2017. The implications of land use/cover dynamics on resources development in Tubah Sub-division, Cameroon. *Journal of the Cameroon Academy of Sciences*, 14(1): 71-85.

Ogoke IJ, Ike GA, Echereobia CO and Ngwuta AA. 2015. Non-destructive leaf area determination in African eggplant (*Solanum macrocarpon*). *Agrosearch*, 15 (2): 13-20.

Ramkhelawan E and Brathwaite RAI. 1990. Leaf area estimation by non-destructive methods in sour orange (*Citrus aurantium* L.). *Tropical Agriculture*, 67(3): 203-206.

Schwarz D and Kalring HP. 2001. Allometry to estimate leaf area of tomato. *Journal of Plant Nutrition*, 24(8): 1291-1309.

Tchinda AT. 2008. *Entandrophragma angolense* (Welw.) C.DC. In: Louppe, D., Oteng-Amoako AA and Brink M (Editors). PROTA (Plant Resources of Tropical Africa/Ressources végétales de l'Afrique tropicale), Wageningen, Netherlands; [cited 2018 September 1]. Available from: <http://prota4u.org/search.asp>.

Ugese FD, Baiyeri KP and Mbah BN. 2008. Leaf area determination of shea butter tree (*Vitellaria paradoxa* C.F. Gaertn.). *International Agrophysics*, 22: 167-170.

[WWF] World Wildlife Fund. 2017. Cameroon Highlands Forests – A Global Ecoregion [cited 2018 August 29]. Available from: clonewwf.wwf-dev.org.

Yuninui NM. 1990. Initiation practical report on Bambui village. A Research Report, Regional College of Agriculture, Bambili, Cameroon.

Zadeh HG, Lorzadeh S and Aryannia N. 2011. Evaluating weeds competitive ability in a corn field in Southern West Iran. *Asian Journal of Crop Science*, 3 (4): 179-187.

Submit your articles online at ecologyresearch.info

Advantages

- Easy online submission
- Complete Peer review
- Affordable Charges
- Quick processing
- Extensive indexing
- You retain your copyright

submit@ecologyresearch.info
www.ecologyresearch.info/Submit.php